arXiv:hep-ph/0702006v1 31 Jan 2007

USM-TH-177

# Flavor separation of the Boer-Mulders function from unpolarized $\pi^- p$ and $\pi^- D$ Drell-Yan processes

Zhun Lu [a], Bo-Qiang Ma [b,*] , Ivan Schmidt [a,*]

[a] *Departamento de Física, Universidad Técnica Federico Santa María, Casilla 110-V, Valparaíso, Chile*

[b] *School of Physics, Peking University, Beijing 100871, China*

---

**Abstract**

We show that measuring the $\cos 2\phi$ angular dependence in unpolarized Drell-Yan processes with $\pi^-$ beams colliding on proton and deuteron targets can determine the ratio of the Boer-Mulders functions for $d$ and $u$ quarks inside the proton $h_1^{\perp,d}/h_1^{\perp,u}$, which is still lack of theoretical constraint. The comparison of the $\cos 2\phi$ asymmetries measured in unpolarized $\pi^- p$ and $\pi^- D$ Drell-Yan processes, which are accessible at CERN by the COMPASS collaboration, can help to discriminate whether $h_1^\perp$ effects or QCD vacuum effects are preferred by data.

*Key words:* Drell-Yan process, azimuthal asymmetry, $\mathbf{k}_T$-dependent distribution functions, transverse spin
*PACS:* 13.85.-t, 12.39.-x, 13.88.+e, 14.20.Dh

---

The transverse spin phenomena appearing in high energy scattering processes [1] is among the most interesting issues of spin physics. Substantial single-spin asymmetries (SSA) in semi-inclusive deeply inelastic scattering (SIDIS) [2,3,4,5], with one incoming nucleon transversely polarized, have been measured by several experiments. The interpretation of these asymmetries provides new insights into QCD and nucleon structure [6,7,8,9,10,11,12]. One of the mechanisms which can account for such SSA is the Sivers effect [6], related to a spin

---

[*] Corresponding authors.
*Email addresses:* `mabq@phy.pku.edu.cn` (Bo-Qiang Ma),
`ivan.schmidt@usm.cl` (Ivan Schmidt).

and $\mathbf{k}_T$-dependent function [13,14,15] named as Sivers function $f_{1T}^{\perp}(x,\mathbf{k}_T^2)$. It arises from the correlation between the nucleon transverse spin and quark transverse momentum. Despite its naively $T$-odd structure, $f_{1T}^{\perp}$ has been shown to be non-zero [10] due to its special gauge-link structure [11,12]. In the case of unpolarized collisions, large $\cos 2\phi$ asymmetries have been observed [16,17] in $\pi^- N$ Drell-Yan dilepton production processes. A parton interpretation of this asymmetry has been proposed in [18] in terms of another leading twist $T$-odd distribution function, the Boer-Mulders function $h_1^{\perp}(x,\mathbf{k}_T^2)$ [9], which has the same QCD origin [11] as the Sivers function. It describes the transverse polarization distribution of the quark inside the unpolarized hadron due to the correlation between the quark transverse spin and transverse momentum. The model calculations of the Boer-Mulders function have been performed in Refs. [19,20,21,22] for the proton and in Ref. [23] for the pion, based on the gauge-link structure of the distribution. The resulting asymmetries have been estimated for unpolarized $p\bar{p}$ [20,24,25] and $\pi^- N$ [26] Drell-Yan processes. The role of Boer-Mulders functions in unpolarized SIDIS process has also been discussed [27,28]. One of the significant features of the Boer-Mulders function relies on that fact that the spin structure of hadrons can be also studied in physical processes without invoking beam or target polarization.

The large anomalous Drell-Yan asymmetry implies a significant non-zero size of the Boer-Mulders function. However its extraction from experimental data is relatively difficult comparing to that of the Sivers function. In the latter case the Sivers function is convoluted with ordinary distribution/fragmentation functions, which are well-known. Several sets of the proton Sivers functions for both $u$ and $d$ quark have been extracted [29,30,31,32] recently from HERMES [3,5] and COMPASS [4] data on single-transverse spin asymmetries in SIDIS. In the process in which the Boer-Mulders function contributes, it is always convoluted with itself or other chiral-odd functions, such as the transversity or the Collins fragmentation function [7], which are also not well known at present. Different models or theoretical considerations [21,22,33] predict very different flavor dependence of $h_1^{\perp}$, and even the relative sign between $h_1^{\perp,u}$ and $h_1^{\perp,d}$ is not known. In this paper we suggest to use unpolarized $\pi^- p$ and $\pi^- D$ Drell-Yan processes to access the flavor dependence of $h_1^{\perp}$, especially the ratio $h_1^{\perp,d}/h_1^{\perp,u}$. The hadron program by the COMPASS collaboration will start in 2007 at CERN, in which a $\pi^-$ beam colliding with both proton and deuteron targets are going to be available. We show that by comparing the $\cos 2\phi$ angular dependent part of the cross sections of unpolarized $\pi^- p$ and $\pi^- D$ Drell-Yan processes, one can determine the ratio $h_1^{\perp,d}/h_1^{\perp,u}$. By comparing the $\cos 2\phi$ asymmetries in these processes, one can also discriminate whether $h_1^{\perp}$ effects or QCD vacuum effects [34] are preferred by data. The later effects can also explain the observed asymmetry in a quite different way. The idea is that the non-perturbative vacuum structure in QCD can lead to fluctuating chromomagnetic vacuum fields. When the $q$ and $\bar{q}$ travel through these fields the spin orientations of the quark pair might be correlated [35] before they annihilate to $\gamma^*$. In Ref. [34] a factorization violating spin correlation is proposed and used to fit the data of the NA10 experiments. According to the color interaction nature of these vacuum effects, the "correlation strengths" for different quark flavors are the same. Therefore vacuum effects imply no flavor dependence of the asymmetries in different processes, which is different from $h_1^\perp$ effects.

The general form of the angular differential cross section for unpolarized Drell-Yan process is

$$\frac{1}{\sigma}\frac{d\sigma}{d\Omega}=\frac{3}{4\pi}\frac{1}{\lambda+3}\left(1+\lambda\cos^2\theta+\mu\sin2\theta\cos\phi+\frac{\nu}{2}\sin^2\theta\cos2\phi\right). \tag{1}$$

The $\cos 2\phi$ angular dependence of the lepton pair can be analyzed in the Collins-Soper frame, in which the unpolarized differential cross section is expressed as

$$\frac{d\sigma(h_A h_B\to l\bar{l}X)}{d\Omega dx_1 dx_2 d^2\mathbf{q}_\perp}=\frac{\alpha^2}{3Q^2}\sum_a e_a^2\Big\{A(y)\mathcal{F}[f_{1a/A}(x_1,\mathbf{p}_\perp^2)f_{1\bar{a}/B}(x_2,\mathbf{k}_\perp^2)]+B(y)\cos2\phi$$
$$\times\mathcal{F}\left[(2\hat{\mathbf{h}}\cdot\mathbf{p}_\perp\hat{\mathbf{h}}\cdot\mathbf{k}_\perp-\mathbf{p}_\perp\cdot\mathbf{k}_\perp)\frac{h_{1a/A}^\perp(x_1,\mathbf{p}_\perp^2)h_{1\bar{a}/B}^\perp(x_2,\mathbf{k}_\perp^2)}{M_A M_B}\right]\Big\}+(a\leftrightarrow\bar{a}), \tag{2}$$

where the notation

$$\mathcal{F}[\cdots]=\int d^2\mathbf{p}_\perp d^2\mathbf{k}_\perp\delta^2(\mathbf{p}_\perp+\mathbf{k}_\perp-\mathbf{q}_\perp)\times\{\cdots\} \tag{3}$$

shows the convolution of transverse momenta, and

$$A(y)=\left(\frac{1}{2}-y+y^2\right)=\frac{1}{4}(1+\cos^2\theta), \tag{4}$$
$$B(y)=y(1-y)=\frac{1}{4}\sin^2\theta \tag{5}$$

in the c.s. frame of the lepton pair. From Eq. (2) we can write down the expression for the $\cos 2\phi$ asymmetry coefficient ($\lambda=1$, $\mu=0$)

$$\nu=\frac{2\mathcal{F}\left[(2\hat{\mathbf{h}}\cdot\mathbf{p}_\perp\hat{\mathbf{h}}\cdot\mathbf{k}_\perp-\mathbf{p}_\perp\cdot\mathbf{k}_\perp)h_{1a/A}^\perp h_{1\bar{a}/B}^\perp\right]}{M_A M_B\mathcal{F}[f_{1a/A}f_{1\bar{a}/B}]}. \tag{6}$$



Fig. 1. $xh_1^{\perp(1),u}(x)$ and $xh_1^{\perp(1),d}(x)$ from different models or theoretical considerations: (a) MIT bag model, (b) spectator model with axial-vector diquark, and (c) large-$N_c$ limit.

The azimuthal angel dependent part can be picked up by a weighting procedure [36]:

$$\left\langle \frac{q_T^2 \cos 2\phi}{4M_A M_B} \right\rangle = \int d\phi d^2 q_T \frac{d\sigma(h_A h_B \to l\bar{l}X)}{d\Omega dx_1 dx_2 d^2\mathbf{q}_\perp} \cdot \frac{q_T^2 \cos 2\phi}{4M_A M_B} \tag{7}$$

$$= \frac{\alpha^2}{3Q^2} B(y) \sum_a e_a^2 h_{1a/A}^{\perp(1)}(x_1) h_{1\bar{a}/B}^{\perp(1)}(x_2) + (a \leftrightarrow \bar{a}), \tag{8}$$

in which

$$h_{1a/H}^{\perp(1)}(x) = \int d^2 k_\perp \frac{k_\perp^2}{2M_H^2} h_{1a/H}^\perp(x, k_\perp^2) \tag{9}$$

is the first $k_\perp^2$-moment of $h_{1a/H}^\perp(x, k_\perp^2)$.

Because of the simple structure of the pion compared to that of the nucleon, it is advantageous to use a pion beam to unravel the quark structure of the nucleon in the Drell-Yan process. In leading twist there are only two distribution functions for the pion, the usual momentum distribution function and the Boer-Mulders function. In Refs. [23,26] calculations based on a quark spectator antiquark model showed that the Boer-Mulders functions of the two valence quarks inside the pion (such as $\bar{u}$ and $d$ inside $\pi^-$) are equal. Actually this relation can be obtained from model independent principles: charge conjugation symmetry and isospin symmetry. In the following we will use $h_{1\pi}^\perp$ and $\bar{h}_{1\pi}^\perp$ to denote the Boer-Mulders functions of the valence and sea quarks inside the pion respectively, and $h_1^{\perp,q(\bar{q})}$ for the Boer-Mulders functions of the quark flavor $q(\bar{q})$ inside the proton. Thus the weighted cross section of the unpolarized $\pi^- p$ Drell-Yan process is ($W = Q_T^2 \cos 2\phi / 4M_A M_B$)

$$\langle W\rangle_{\pi^- p}(x_1,x_2) = \frac{\alpha^2}{3Q^2}B(y)\left\{h_{1\pi}^{\perp(1)}(x_1)\left[e_u^2 h_1^{\perp(1),u}(x_2)+e_d^2 h_1^{\perp(1),\bar{d}}(x_2))\right]\right.$$
$$\left.+\bar{h}_{1\pi}^{\perp(1)}(x_1)\left[e_u^2 h_1^{\perp(1),\bar{u}}(x_2)+e_d^2 h_1^{\perp(1),d}(x_2))\right]\right\}. \quad (10)$$

We are interested in the region were $x_{1/2}$ are not so small, where the sea quark contribution to the $\cos 2\phi$ angular dependent cross section is expected to be small. Then we can use the approximation

$$\langle W\rangle_{\pi^- p}(x_1,x_2) \approx \frac{4\alpha^2}{3Q^2}B(y)e_u^2 h_{1\pi}^{\perp(1)}(x_1)h_1^{\perp(1),u}(x_2). \quad (11)$$

The cross section for the $\pi^- D$ Drell-Yan process can be approximated as $\sigma_{\pi^- D} = \sigma_{\pi^- p} + \sigma_{\pi^- n}$, thus

$$\langle W\rangle_{\pi^- D}(x_1,x_2) = \langle W\rangle_{\pi^- p}(x_1,x_2) + \langle W\rangle_{\pi^- n}(x_1,x_2). \quad (12)$$

The Boer-Mulders functions of quarks inside the neutron can be related to those in the proton: $h_{1u/n}^{\perp} = h_1^{\perp,d}$, $h_{1d/n}^{\perp} = h_1^{\perp,u}$. Therefore

$$\langle W\rangle_{\pi^- n}(x_1,x_2) = \frac{\alpha^2}{3Q^2}B(y)\left\{h_{1\pi}^{\perp(1)}(x_1)\left[e_u^2 h_1^{\perp(1),d}(x_2)+e_d^2 h_1^{\perp(1),\bar{u}}(x_2))\right]\right.$$
$$\left.+\bar{h}_{1\pi}^{\perp(1)}(x_1)\left[e_u^2 h_1^{\perp(1),\bar{d}}(x_2)+e_d^2 h_1^{\perp(1),u}(x_2))\right]\right\}. \quad (13)$$

Ignoring the sea quark contribution, we have

$$\langle W\rangle_{\pi^- D}(x_1,x_2) \approx \frac{\alpha^2}{3Q^2}B(y)e_u^2 h_{1\pi}^{\perp(1)}(x_1)(h_1^{\perp(1),u}(x_2)+h_1^{\perp(1),d}(x_2)). \quad (14)$$

Then the ratio of the weighted cross sections for unpolarized $\pi^- p$ and $\pi^- D$ Drell-Yan processes is

$$\frac{\langle W\rangle_{\pi^- D}(x_1,x_2)}{2\langle W\rangle_{\pi^- p}(x_1,x_2)} = \frac{1}{2}\left(1+\frac{h_1^{\perp(1),d}(x_2)}{h_1^{\perp(1),u}(x_2)}\right). \quad (15)$$

The Drell-Yan process at COMPASS can explore the kinematics region $s = 400\,\mathrm{GeV}^2$ and $Q^2 = 20\,\mathrm{GeV}^2$, where $x_1x_2 = Q^2/s = 0.05$. Therefore the Drell-Yan experiments at COMPASS can access the quark structure in the valence regime $0.1 \leq x_{1/2} \leq 0.5$, where Eq. (15) is a good approximation. Then the ratio of the Boer-Mulders functions of $d$ and $u$ quarks is

$$\frac{h_1^{\perp(1),d}(x_2)}{h_1^{\perp(1),u}(x_2)} = \frac{\langle W\rangle_{\pi^- D}(x_1,x_2)}{\langle W\rangle_{\pi^- p}(x_1,x_2)} - 1. \tag{16}$$

Another possibility is to use a $\pi^+$ beam to collide on the proton, which process is sensitive to the $d$ quark distribution of the proton:

$$\frac{\langle W\rangle_{\pi^+ p}(x_1,x_2)}{\langle W\rangle_{\pi^- p}(x_1,x_2)} = \frac{h_1^{\perp(1),d}(x_2)}{4h_1^{\perp(1),u}(x_2)}. \tag{17}$$

This relation can serve as cross check for Eq. (16).

knowledge of the Boer-Mulders functions is very poor, including its size and shape, and the relative size between $h_1^{\perp,u}$ and $h_1^{\perp,d}$. In the last case different models or theoretical considerations predict quite different values for the ratio $h_1^{\perp,d}/h_1^{\perp,u}$. A calculation based on the MIT bag model gave [21] $h_1^{\perp,d} = \frac{1}{2}h_1^{\perp,u}$. The spectator model with axial-vector diquarks predicted [22] that $h_1^{\perp,d}$ has a opposite sign and a rather larger size compared with $h_1^{\perp,u}$, while large-$N_c$ showed that $h_1^{\perp,d} = h_1^{\perp,u}$ [33] modulo $1/N_c$ corrections. In Fig. 1 we show the curves for $h_1^{\perp(1),u}(x)$ and $h_1^{\perp(1),d}(x)$ from these models or theoretical considerations. In the case of large-$N_c$ since there is no detail calculation on the size of $h_1^{\perp}$, we just naively assume $h_1^{\perp(1),u} = f_{1T}^{\perp(1),u}$ (these two function are closely related and expected to have similar size), and for $f_{1T}^{\perp(1),u}$ we adopt the parametrization [31] extracted from HERMES experimental data based on large-$N_c$. In the case of MIT bag model the ratio in Eq. (15) is 0.75 and in large-$N_c$ consideration is 1, while in axial-vector diquark model the ratio is mostly negative. These predictions can be tested by measuring the ratio $\langle W\rangle_{\pi^- D}/(2\langle W\rangle_{\pi^- p})$. Vice versa, according to Eq. (16), from the measured ratio $\langle W\rangle_{\pi^- D}/(2\langle W\rangle_{\pi^- p})$ the size of $h_1^{\perp,d}/h_1^{\perp,u}$ can be determined.

The analysis can be easily extended to the single (transverse) polarized Drell-Yan processes $\pi p^\uparrow(D^\uparrow) \to \mu^+\mu^- X$ at COMPASS. The measurement of the single-spin asymmetries in this process is also important for the more precise extraction of the Sivers function (through $f_{1\pi} \otimes f_{1T}^{\perp}$), and to test the QCD prediction $f_{1T}^{\perp}|_{\mathrm{SIDIS}} = -f_{1T}^{\perp}|_{\mathrm{DY}}$ [11,37], and can provide alternative access to transversity (through $h_{1\pi}^{\perp} \otimes h_1$) [18]. In single polarized Drell-Yan processes the relations similar to Eqs. (15) and (17) can be deduced for the Sivers function and for transversity.

It is also interesting to compare the $\cos 2\phi$ asymmetries in unpolarized $\pi^- p$ and $\pi^- D$ processes. The anomalous Drell-Yan asymmetry might be produced by $h_1^{\perp}$, which is labelled as hadronic effect. Nevertheless, there is another effect that could also account for this asymmetry, the so called QCD vacuum effect [34]. A significant difference between these two effects [38] is that the



Fig. 2. The weighted $\cos 2\phi$ asymmetries in unpolarized $\pi^- p$ (solid line) and $\pi^- D$ (dashed line) Drell-Yan processes. (a) The MIT Bag model result. The two asymmetries are equal since $f_1^d/f_1^u = 1/2$ in the lowest bag mode in which case is used in calculation. (b) The axial-vector diquark model result. (c) The result from large-$N_c$.

first one is flavor dependent, while the latter is flavor blind. Thus comparing the asymmetries in different processes can discriminate which effect is really responsible for the asymmetry. In the case of the hadronic effect, the weighted $\cos 2\phi$ asymmetries of these two processes can be calculated from following expressions:

$$\hat{\nu}_p = a_{UU} \frac{\langle W \rangle_{\pi^- p}}{\langle 1 \rangle_{\pi^- p}} = \frac{h_{1\pi}^{\perp(1)}(x_1) h_1^{\perp(1),u}(x_2)}{f_{1\pi}(x_1) f_1^u(x_2)}, \tag{18}$$

$$\hat{\nu}_D = a_{UU} \frac{\langle W \rangle_{\pi^- D}}{\langle 1 \rangle_{\pi^- D}} = \frac{h_{1\pi}^{\perp(1)}(x_1)(h_1^{\perp(1),u}(x_2) + h_1^{\perp(1),d}(x_2))}{f_{1\pi}(x_1)(f_1^u(x_2) + f_1^d(x_2))}, \tag{19}$$

where $a_{UU} = (1 + \cos^2\theta)/\sin^2\theta$, and $\langle 1 \rangle$ is calculated from Eq. (7) when $W$ is replaced by 1. Again we neglect the sea contribution here. The ratio of the asymmetries in these two processes is then

$$\frac{\hat{\nu}_D}{\hat{\nu}_p} = \frac{1 + \frac{h_1^{\perp(1),d}(x_2)}{h_1^{\perp(1),u}(x_2)}}{1 + \frac{f_1^d(x_2)}{f_1^u(x_2)}}. \tag{20}$$

The QCD vacuum effect will yield $\hat{\nu}_p = \hat{\nu}_D$. In Fig. 2 we show the asymmetries calculated from Eq. (18) and (19) using the three options of Boer-Mulders functions given in Fig. 1. For the unpolarized distributions, which appear in the denominators in Eq. (18) and (19) we adopt well known model results. We do not aim at a precise estimate of the asymmetries, only to a rough comparison among them. In the case that $h_1^{\perp(1),u}/f_1^u$ is different from $h_1^{\perp(1),d}/f_1^d$ (equivalently $h_1^{\perp(1),d}/h_1^{\perp(1),u}$ different from $f_1^d/f_1^u$), different asymmetries will be observed in these two processes, as shown in Fig. 2b and Fig. 2c. Therefore the difference appearing in the $\cos 2\phi$ asymmetries in unpolarized $\pi^- p$ and $\pi^- D$ Drell-Yan processes can provide evidence whether hadronic effects or QCD vacuum effects are preferred by data.

In summary, we have showed that by comparing the $\cos 2\phi$ angular dependent part of the cross sections of the unpolarized $\pi^- p$ and $\pi^- D$ Drell-Yan processes, which will be accessible in future COMPASS experiments, one can determine the ratio $h_1^{\perp,d}/h_1^{\perp,u}$ in the valence region, which still lacks theoretical input. By checking the difference of the $\cos 2\phi$ asymmetries in these two processes, one can also provide a discrimination whether hadronic effects or QCD vacuum effects are preferred by data. The investigation will lead to better understanding about the role of the quark transverse motion and transverse spin in hadronic reactions.

**Acknowledgements.** This work is partially supported by National Natural Science Foundation of China (Nos. 10421503, 10575003, 10505001, 10528510), by the Key Grant Project of Chinese Ministry of Education (No. 305001), by the Research Fund for the Doctoral Program of Higher Education (China), by Fondecyt (Chile) under Project No. 3050047 and No. 1030355.

## References

[1] For a review on tranverse polarization phenomena, see V. Barone, A. Drago, P.G. Ratcliffe, Phys. Rep. 359 1 (2002).

[2] A. Bravar et al., SMC Collaboration, Nucl. Phys. A 666 (2000) 314.

[3] A. Airapetian et al., HERMES Collaboration, Phys. Rev. Lett. 94 (2005) 012002.

[4] V.Yu. Alexakhin et al., COMPASS Collaboration, Phys. Rev. Lett. 94 (2005) 202002.

[5] M. Diefenthaler, HERMES Collaboration, in Proceedings of DIS 2005, Madison, Wisconsin (USA), hep-ex/0507013.

[6] D. Sivers, Phys. Rev. D 41 (1990) 83;
D. Sivers, Phys. Rev. D 43 (1991) 261.

[7] J.C. Collins, Nucl. Phys. B 396 (1993) 161.

[8] M. Anselmino, M. Boglione, F. Murgia, Phys. Lett. B 362 164 (1995).
M. Anselmino, F. Murgia, Phys. Lett. B 442, 470 (1998).

[9] D. Boer, P.J. Mulders, Phys. Rev. D 57 (1998) 5780.

[10] S.J. Brodsky, D.S. Hwang, I. Schmidt, Phys. Lett. B 530 (2002) 99;
S.J. Brodsky, D.S. Hwang, I. Schmidt, Nucl. Phys. B 642 (2002) 344.

[11] J.C. Collins, Phys. Lett. B 536 (2002) 43.

[12] X. Ji and F. Yuan, Phys. Lett. B 543 (2002) 66;
A.V. Belitsky, X. Ji, F. Yuan, Nucl. Phys. B 656 (2003) 165;
D. Boer, P.J. Mulders, F. Pijlman, Nucl. Phys. B 667 (2003) 201.

[13] J. Levelt and P.J. Mulders, Phys. Rev. D 49 (1994) 96.

[14] A. Kotzinian, Nucl. Phys. B 441 (1995) 234.

[15] P.J. Mulders and R.D. Tangerman, Nucl. Phys. B 461 (1996)

[16] S. Falciano et al. NA10 Collaboration, Z. Phys. C 31 (1986) 513;
M. Guanziroli et al. NA10 Collaboration, Z. Phys. C 37 (1988) 545.

[17] J.S. Conway et al. Phys. Rev. D 39 (1989) 92.

[18] D. Boer, Phys. Rev. D 60 (1999) 014012.

[19] G.R. Goldstein, L. Gamberg, Talk given at 31st International Conference on High Energy Physics (ICHEP 2002), Amsterdam, The Netherlands, 24-31 July 2002, hep-ph/0209085.

[20] D. Boer, S.J. Brodsky, D.S. Hwang, Phys. Rev. D 67 (2003) 054003.

[21] F. Yuan, Phys. Lett. B 575 (2003) 45.

[22] A. Bacchetta, A. Schäfer, J.-J. Yang, Phys. Lett. B 578 (2004) 109.

[23] Z. Lu, B.-Q. Ma, Phys. Rev. D 70 (2004) 094044.

[24] G. R. Goldstein, L. Gamberg, hep-ph/0506127.

[25] V. Barone, Z. Lu, B.-Q. Ma, hep-ph/0612350, to appear in European Physical Journal C.

[26] Z. Lu, B.-Q. Ma, Phys. Lett. B 615 (2005) 200.

[27] L.P. Gamberg, G.R. Goldstein, K.A. Oganessyan, Phys. Rev. D 67 (2003) 071504;
L.P. Gamberg, G.R. Goldstein , K.A. Oganessyan, Phys. Rev. D 68 (2003) 051501(R).

[28] V. Barone, Z. Lu, B.-Q. Ma, Phys. Lett. B 632 (2006) 277.

[29] M. Anselmino, M. Boglione, U. D'lesio, A. Kotzinian, F. Murgia, A. Prokudin, Phys. Rev. D 72 (2005) 094007;
M. Anselmino, M. Boglione, U. D'lesio, A. Kotzinian, F. Murgia, A. Prokudin, Phys. Rev. D 72 (2005) 099903, Erratum.

[30] W. Vogelsang , F. Yuan, Phys. Rev. D 72 (2005) 054028.

[31] J.C. Collins, A.V. Efremov, K. Goeke, S. Menzel, A. Metz, P. Schweitzer, Phys. Rev. D73 (2006) 014021;
J.C. Collins, A. V. Efremov, K. Goeke, M. Grosse Perdekamp, S. Menzel, B. Meredith, A. Metz, P. Schweitzer, hep-ph/0510342.

[32] For a comparison to this extraction, see M. Anselmino et al., hep-ph/0511017.

[33] P.V. Pobylitsa, hep-ph/0301236.

[34] A. Brandenburg, O. Nachtmann, E. Mirkes, Z. Phys. C 60 (1993) 697.

[35] O. Nachtmann, A. Reiter, Z. Phys. C 24 (1984) 283.

[36] A. Kotzinian, P.J. Mulders, Phys. Letts. B 406 (1997) 373.

[37] A.V. Efremov, K. Goeke, S. Menzel, A. Metz, P. Schweitzer, Phys. Lett. B 612 (2005) 233.

[38] D. Boer, A. Brandenburg, O. Nachtmann, Utermann, Eur. Phys. J. C 40 (05) 55.